Preprint, `arXiv:1210:5237`

DETERMINING x OR y MOD p^2 WITH $p = x^2 + dy^2$

Zhi-Wei Sun

Department of Mathematics, Nanjing University
Nanjing 210093, People's Republic of China
zwsun@nju.edu.cn
`http://math.nju.edu.cn/∼zwsun`

arXiv:1210.5237v3 [math.NT] 25 Oct 2012

Abstract. Let p be an odd prime and let $d \in \{2,3,7\}$. When $(\frac{-d}{p}) = 1$ we can write $p = x^2 + dy^2$ with $x, y \in \mathbb{Z}$; in this paper we aim at determining x or y modulo p^2. For example, when $p = x^2 + 3y^2$, we show that if $p \equiv x \equiv 1 \pmod 4$ then

$$\sum_{k=0}^{(p-1)/2} (3[3 \mid k] - 1)(2k+1)\frac{\binom{2k}{k}^2}{(-16)^k} \equiv \left(\frac{2}{p}\right) 2x \pmod{p^2}$$

where $[3 \mid k]$ takes 1 or 0 according as $3 \mid k$ or not, and that if $-p \equiv y \equiv 1 \pmod 4$ then

$$\sum_{k=0}^{(p-1)/2} \left(\frac{k}{3}\right) \frac{k\binom{2k}{k}^2}{(-16)^k} \equiv (-1)^{(p+1)/4} y \equiv \sum_{k=0}^{(p-1)/2} (1 - 3[3 \mid k])\frac{k\binom{2k}{k}^2}{(-16)^k} \pmod{p^2}.$$

1. Introduction

Let $p \equiv 1 \pmod 4$ be a prime. It is well-known that we may write $p = x^2 + y^2$ with $x \equiv 1 \pmod 4$ and $y \equiv 0 \pmod 2$. In 1828 Gauss determined x mod p via the congruence

$$\binom{(p-1)/2}{(p-1)/4} \equiv 2x \pmod p$$

(see, e.g., [BEW, Chapter 9]). In 1986, S. Chowla, B. Dwork and R. J. Evans [CDE] showed further that

$$\binom{(p-1)/2}{(p-1)/4} \equiv \frac{2^{p-1}+1}{2}\left(2x - \frac{p}{2x}\right) \pmod{p^2},$$

2010 *Mathematics Subject Classification.* Primary 11E25, 11A07; Secondary 05A10, 11B65.
Keywords. Representations of primes by binary quadratic forms., congruences
Supported by the National Natural Science Foundation (grant 11171140) of China.

which implies the congruence

$$\binom{(p-1)/2}{(p-1)/4}^2 \equiv 2^{p-1}(4x^2-2p) \pmod{p^2}.$$

Let p be an odd prime. In 1997 van Hamme [vH] showed that

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{64^k} \equiv \begin{cases} 4x^2-2p \pmod{p^2} & \text{if } p\equiv 1 \pmod 4 \ \&\ p=x^2+y^2\ (2\nmid x), \\ 0 \pmod{p^2} & \text{if } p\equiv 3 \pmod 4. \end{cases} \tag{1.0}$$

By Stirling's formula, $n! \sim \sqrt{2\pi n}(n/e)^n$ and hence

$$\lim_{k\to\infty}\sqrt[k]{\binom{2k}{k}^3}=64.$$

Thus we call $\sum_{k=0}^{p-1}\binom{2k}{k}^3/m^k$ mod p^2 the *border case* if $m=64$. In 2009 the author posed several conjectures on $\sum_{k=0}^{p-1}\binom{2k}{k}^3/m^k$ mod p^2 in non-border cases. For example, in Nov. 2009 he conjectured that

$$\sum_{k=0}^{p-1}\binom{2k}{k}^3 \equiv \begin{cases} 4x^2-2p \pmod{p^2} & \text{if } (\frac{p}{7})=1 \ \&\ p=x^2+7y^2, \\ 0 \pmod{p^2} & \text{if } (\frac{p}{7})=-1. \end{cases} \tag{1.1}$$

This appeared as Conjecture 5.3 of [Su1]. In [Su2, Conjecture 5.2] the author also formulated conjectures on $\sum_{k=0}^{p-1}\binom{2k}{k}^3/m^k$ with $m=-8,16,-64,256,-512,4096$. Namely,

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{(-8)^k} \equiv \sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{(-512)^k} \equiv \sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{64^k} \ \left(\mathrm{mod}\ p^{(5+(\frac{-1}{p})/2)}\right), \tag{1.2}$$

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{(-64)^k} \equiv \begin{cases} 4x^2-2p \pmod{p^2} & \text{if } p=x^2+2y^2, \\ 0 \pmod{p^2} & \text{if } p\equiv 5,7 \pmod 8, \end{cases} \tag{1.3}$$

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{16^k} \equiv \begin{cases} 4x^2-2p \pmod{p^2} & \text{if } p=x^2+3y^2, \\ 0 \pmod{p^2} & \text{if } p\equiv 2 \pmod 3, \end{cases} \tag{1.4}$$

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{256^k} \equiv \left(\frac{-1}{p}\right)\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{16^k} \ \left(\mathrm{mod}\ p^{(5+(\frac{p}{3}))/2}\right), \tag{1.5}$$

and

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{4096^k} \equiv \left(\frac{-1}{p}\right)\sum_{k=0}^{p-1}\binom{2k}{k}^3 \ \left(\mathrm{mod}\ p^{(5+(\frac{p}{7}))/2}\right). \tag{1.6}$$

Note that [A, Theorem 5] implies (1.1)-(1.6) modulo p. Z.-H. Sun [S2] got partial success in proving the above conjectures, he showed that

$$\begin{gathered}\sum_{k=0}^{p-1}\binom{2k}{k}^3 \equiv \sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{4096^k} \equiv 0 \pmod{p^2} \ \text{ if } \left(\frac{p}{7}\right) = -1,\\ \sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{(-8)^k} \equiv \sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{(-512)^k} \equiv 0 \pmod{p^2} \ \text{ if } p \equiv 3 \pmod 4,\\ \sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{16^k} \equiv \sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{256^k} \equiv 0 \pmod{p^2} \ \text{ if } p \equiv 2 \pmod 3,\end{gathered}$$

and

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{(-8)^k} \equiv 4x^2 - 2p \pmod{p^2} \quad \text{if } p = x^2 + y^2 \ (2 \nmid x).$$

The key tool of Z.-H. Sun [S2] is the following identity

$$P_n(x)^2 = \sum_{k=0}^{n}\binom{n}{k}\binom{n+k}{k}\binom{2k}{k}\left(\frac{x^2-1}{4}\right)^k, \tag{1.7}$$

where $P_n(x)$ is the Legendre polynomial of degree n given by

$$P_n(x) = \sum_{k=0}^{n}\binom{n}{k}\binom{n+k}{k}\left(\frac{x-1}{2}\right)^k.$$

Note that (1.7) follows from the well-known Clausen identity

$${}_2F_1\left(\begin{matrix}2a, 2b\\ a+b+1/2\end{matrix}\middle| z\right)^2 = {}_3F_2\left(\begin{matrix}2a, 2b, a+b\\ a+b+1/2, 2a+2b\end{matrix}\middle| 4z(1-z)\right)$$

in the special case $a = -n/2$, $b = (n+1)/2$ and $z = (1-x)/2$.

Concerning (1.4) we mention that the author [Su3] proved that

$$\sum_{k=0}^{p-1}(-1)^k A_k \equiv \sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{16^k} \pmod{p^2} \tag{1.8}$$

for any odd prime p, where $A_0, A_1, A_2, \ldots$ are Apéry numbers given by

$$A_n = \sum_{k=0}^{n}\binom{n}{k}^2\binom{n+k}{k}^2 \ (n = 0, 1, 2, \ldots).$$

During their initial attempt to prove (1.1) in 2010, M. Jameson and K. Ono [JO] realized that

$$\sum_{k=0}^{(p-1)/2}\binom{2k}{k}^3(H_{2k}-H_k)\equiv 0\pmod p\quad\text{for any prime } p>3$$

but they did not have a proof of this observation, where H_k denotes the harmonic number $\sum_{0<j\leqslant k}1/j$. When $p>3$ is a prime with $p\equiv 3\pmod 4$, by [Su1, (1.11)] and (1.0) we have

$$\sum_{k=0}^{(p-1)/2}\frac{\binom{2k}{k}^3}{64^k}H_k\equiv 0\ (\mathrm{mod}\ p).$$

On March 2, 2010 the author formulated a conjecture (which was presented as [Su, Conj. A37]) on $\sum_{k=0}^{p-1}\binom{2k}{k}^3(H_{2k}-H_k)/m^k$ modulo p or p^2, where $p>3$ is a prime and $m\in\{1,-8,16,-64,256,-512,4096\}$.

Quite recently the author's conjectural congruences (1.1)-(1.6) modulo p^2 were confirmed by J. Kibelbek, L. Long, K. Moss, B. Sheller and H. Yuan [KLMSY]. They also proved for any prime $p>3$ and $m=1,-8,16,-64,256,-512,4096$ the congruence

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{m^k}(H_{2k}-H_k)\equiv\frac{q_p(m)}{6}\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{m^k}\ (\mathrm{mod}\ p)\tag{1.9}$$

where $q_p(m)$ denotes the Fermat quotient $(m^{p-1}-1)/p$.

The work of [KLMSY] depends heavily on the paper [CH] of M. J. Coster and L. van Hamme [CV] in which the authors determined

$$P_n(\sqrt2),\ P_n\left(\frac{3\sqrt2}{4}\right),\ P_n(\sqrt{-3}),\ P_n\left(\frac{\sqrt3}{2}\right),\ P_n\left(3\sqrt{-7}\right),\ P_n\left(\frac{3\sqrt7}{8}\right)$$

modulo p^2 via complex multiplication of elliptic curves, where $p=2n+1$ is an odd prime.

When $p=x^2+y^2$ with $4\mid x-1$ and $2\mid y$, the author [Su5] showed that we can determine x mod p^2 in the following way:

$$\left(\frac{2}{p}\right)x\equiv\sum_{k=0}^{(p-1)/2}\frac{k+1}{8^k}\binom{2k}{k}^2\equiv\sum_{k=0}^{(p-1)/2}\frac{2k+1}{(-16)^k}\binom{2k}{k}^2\pmod{p^2}.$$

With the help of Coster-van Hamme's work, in this paper we determine x or y mod p^2 when $p=x^2+dy^2$ with $d=2,3,7$.

Before stating our main results we need to introduce Lucas sequences.

Let $A, B \in \mathbb{Z}$. The Lucas sequence $u_n = u_n(A, B)$ $(n \in \mathbb{N})$ and its companion $v_n = v_n(A, B)$ $(n \in \mathbb{N})$ are defined as follows:

$$\begin{aligned} u_0 = 0,\ u_1 = 1,\ &\text{and } u_{n+1} = Au_n - Bu_{n-1}\ (n = 1, 2, 3, \dots);\\ v_0 = 2, v_1 = A,\ &\text{and } v_{n+1} = Av_n - Bv_{n-1}\ (n = 1, 2, 3, \dots). \end{aligned}$$

It is well known that $(\alpha - \beta)u_n = \alpha^n - \beta^n$ and $v_n = \alpha^n + \beta^n$ for all $n \in \mathbb{N}$, where α and β are the two roots of the equation $x^2 - Ax + B = 0$. The sequence $P_n = u_n(2, -1)$ $(n \in \mathbb{N})$ is called the Pell sequence and its companion is the sequence $Q_n = v_n(2, 1)$ $(n \in \mathbb{N})$.

Theorem 1.1. *Let p be a prime with $p \equiv 1, 3 \pmod 8$. And write $p = x^2 + 2y^2$ with $x, y \in \mathbb{Z}$ and $x \equiv 1 \pmod 4$.*

(i) *If $p \equiv 1 \pmod 8$, then*

$$\begin{aligned} 4\sum_{k=0}^{p-1}\binom{2k}{k}^2\frac{kP_k}{32^k} \equiv &\sum_{k=0}^{p-1}\binom{2k}{k}^2\frac{kQ_k}{32^k}\\ \equiv &(-1)^{(p-1)/8+(x-1)/4}\left(\frac{p}{x} - 2x\right) \pmod{p^2}, \end{aligned}$$

and we can determine x mod p^2 via the congruence

$$(-1)^{(x-1)/4}x \equiv \frac{(-1)^{(p-1)/8}}{2}\sum_{k=0}^{p-1}\binom{2k}{k}^2\frac{(k+1)Q_k}{32^k} \pmod{p^2}.$$

(ii) *If $p \equiv 3 \pmod 8$, then*

$$\sum_{k=0}^{p-1}\binom{2k}{k}^2\frac{kP_k}{32^k} \equiv \frac{1}{2}\sum_{k=0}^{p-1}\binom{2k}{k}^2\frac{kQ_k}{32^k} \equiv (-1)^{(y+1)/2}y \pmod{p^2}$$

and

$$\sum_{k=0}^{p-1}\binom{2k}{k}^2\frac{P_k}{32^k} \equiv (-1)^{(y-1)/2}\left(2y - \frac{p}{4y}\right) \pmod{p^2}.$$

Remark 1.1. Theorem 1.1 was first conjectured by the author in 2009, it appeared as part of [Su4, Conjecture 2.3].

Theorem 1.2. *Let $p \equiv 1 \pmod 3$ be a prime and write $p = x^2 + 3y^2$ with $x, y \in \mathbb{Z}$.*

(i) *If $p \equiv x \equiv 1 \pmod 4$, then we can determine x mod p^2 via the congruence*

$$\left(\frac{2}{p}\right)2x \equiv \sum_{k=0}^{p-1}(3[3 \mid k] - 1)(2k + 1)\frac{\binom{2k}{k}^2}{(-16)^k} \pmod{p^2}. \tag{1.10}$$

(ii) *If* $-p\equiv y\equiv 1\pmod 4$*, then*

$$\sum_{k=0}^{p-1}\left(\frac{k}{3}\right)\frac{\binom{2k}{k}^2}{(-16)^k}\equiv(-1)^{(p-3)/4}\left(4y-\frac{p}{3y}\right)\pmod{p^2},\tag{1.11}$$

and we can determine y *mod* p^2 *in the following way:*

$$\sum_{k=0}^{p-1}\left(\frac{k}{3}\right)\frac{k\binom{2k}{k}^3}{(-16)^k}\equiv(-1)^{(p+1)/4}y\equiv\sum_{k=0}^{p-1}(1-3[3\mid k])\frac{k\binom{2k}{k}^2}{(-16)^k}\pmod{p^2}.\tag{1.12}$$

Remark 1.2. Part (ii) of Theorem 1.2 appeared as part (i) of [Su2, Conjecture 5.11].

Theorem 1.3. *Let* p *be an odd prime with* $(\frac{p}{7})=1$ *and write* $p=x^2+7y^2$ *with* $x,y\in\mathbb{Z}$*.*

(i) *If* $p\equiv x\equiv 1\pmod 4$*, then we may determine* x *mod* p^2 *via the congruence*

$$\sum_{k=0}^{p-1}(4k+3)\frac{\binom{2k}{k}^2}{16^k}v_k(1,16)\equiv 6\left(\frac{2}{p}\right)x\pmod{p^2}.\tag{1.13}$$

(ii) *If* $-p\equiv y\equiv 1\pmod 4$*, then we may determine* y *mod* p^2 *in the following way:*

$$\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^2}{16^k}u_k(1,16)\equiv\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^2}{16^k}v_k(1,16)\equiv-\left(\frac{2}{p}\right)\frac{y}{2}\pmod{p^2}.\tag{1.14}$$

Theorem 1.4. *Let* $p>3$ *be a prime.*

(i) *If* $p\equiv 1\pmod{12}$ *and* $p=x^2+3y^2$ *with* $x,y\in\mathbb{Z}$ *and* $x\equiv 1\pmod 4$*, then*

$$\sum_{k=0}^{p-1}\binom{2k}{k}^2\frac{v_k(4,1)}{64^k}\equiv 4x-\frac{p}{x}\ (\mathrm{mod}\ p^2)$$

and we can determine x *mod* p^2 *by*

$$\sum_{k=0}^{p-1}\binom{2k}{k}^2\frac{(k-1)v_k(4,1)}{64^k}\equiv -2x\ (\mathrm{mod}\ p^2).$$

(ii) *If* $p\equiv 7\pmod{12}$ *and* $p=x^2+3y^2$ *with* $x,y\in\mathbb{Z}$ *and* $y\equiv 1\pmod 4$*, then*

$$\sum_{k=0}^{p-1}\frac{u_k(4,1)}{64^k}\binom{2k}{k}^2\equiv 2y-\frac{p}{6y}\ (\mathrm{mod}\ p^2)$$

and also

$$y\equiv\sum_{k=0}^{p-1}\binom{2k}{k}^2\frac{ku_k(4,1)}{64^k}\equiv\frac{1}{4}\sum_{k=0}^{p-1}\binom{2k}{k}^2\frac{kv_k(4,1)}{64^k}\ (\mathrm{mod}\ p^2).$$

Remark 1.3. Theorem 1.4 was part of [Su4, Conjecture 2.4].

In contrast with (1.9), we have the following theorem.

Theorem 1.5. *Let p be an odd prime and let $a, b, m \in \mathbb{Z}$ with $m \not\equiv 0 \pmod p$. Then*

$$\begin{aligned}&\left(\frac{-m}{p}\right)\sum_{k=0}^{p-1}\frac{ak+b}{m^k}\binom{2k}{k}^3+\frac{\bar{m}^{p-1}+1}{4}\sum_{k=0}^{p-1}\frac{2ak+a-2b}{\bar{m}^k}\binom{2k}{k}^3\\ \equiv&\frac{ap}{2}\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{\bar{m}^k}+\frac{3p}{2}\sum_{k=0}^{p-1}\frac{2ak+a-2b}{\bar{m}^k}\binom{2k}{k}^3(H_{2k}-H_k) \pmod{p^2},\end{aligned}\tag{1.15}$$

where $\bar{m} = 2^{12}/m$.

Clearly Theorem 1.5 with $a = 0$ and $b = 1$ yields the following corollary.

Corollary 1.1. *Let p be an odd prime and let $m \in \mathbb{Z}$ with $p \nmid m$. Set $\bar{m} = 4096/m$. Then*

$$\begin{aligned}&\left(\frac{-m}{p}\right)\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{m^k}-\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{\bar{m}^k}\\ \equiv&-\frac{p}{2}\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{\bar{m}^k}(6(H_{2k}-H_k)-q_p(\bar{m})) \pmod{p^2}.\end{aligned}$$

Remark 1.4. In view of Corollary 1.1, for any odd prime $p > 3$, (1.9) holds for all $m \in \{1, -8, 16, -64, 256, -512, 1024\}$, if and only if the symmetric relation

$$\left(\frac{-m}{p}\right)\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{m^k}\equiv\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{\bar{m}^k} \pmod{p^2}\tag{1.16}$$

holds for all $m \in \{1, -8, 16, -64, 256, -512, 1024\}$. The latter was conjectured by the author in [Su2, Conj. 5.2].

In the next section shall provide some lemmas. We are going to show Theorems 1.1-1.4 in Section 3. In Section 4 we will prove Theorem 1.5 and derive several corollaries of Th. 1.5. Section 4 also contains a conjecture involving harmonic numbers.

2. Several lemmas

Lemma 2.1. *Let p be an odd prime and let $m \in \mathbb{Z}$ with $m \not\equiv 0 \pmod p$. Let m_* be a root of the equation $x^2 - mx + 16m = 0$ in the ring of algebraic p-adic integers. Then, for any algebraic p-adic integers a and b, we have*

$$\begin{aligned}&\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{m^k}\left(\frac{ak}{16}(m_*-m+32)+b\right)\\ \equiv&2a\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^2}{m_*^k}\sum_{j=0}^{p-1}\frac{\binom{2j}{j}^2}{m_*^j}+b\bigg(\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{m_*^k}\bigg)^2 \pmod{p^2}.\end{aligned}\tag{2.1}$$

Proof. Set

$$S:=\sum_{n=0}^{p-1}\frac{an+b}{m_*^n}\sum_{k=0}^{n}\binom{2k}{k}^2\binom{2(n-k)}{n-k}^2.$$

By [Su5, Lemma 3.1],

$$\begin{aligned}S=&\sum_{n=0}^{p-1}\frac{an+b}{m_*^n}\sum_{k=0}^{n}\binom{2k}{k}^3\binom{k}{n-k}(-16)^{n-k}\\=&\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^3}{m_*^k}\sum_{n=k}^{p-1}(an+b)\binom{k}{n-k}\left(-\frac{16}{m_*}\right)^{n-k}\\\equiv&\sum_{k=0}^{(p-1)/2}\frac{\binom{2k}{k}^3}{m_*^k}\sum_{j=0}^{k}(a(k+j)+b)\binom{k}{j}\left(-\frac{16}{m_*}\right)^{j}\pmod{p^2}.\end{aligned}$$

Thus

$$\begin{aligned}S\equiv&\sum_{k=0}^{(p-1)/2}\frac{\binom{2k}{k}^3}{m_*^k}\left((ak+b)\left(1-\frac{16}{m_*}\right)^k+ak\sum_{0<j\leqslant k}\binom{k-1}{j-1}\left(-\frac{16}{m_*}\right)^j\right)\\=&\sum_{k=0}^{(p-1)/2}\frac{\binom{2k}{k}^3}{m_*^k}\left((ak+b)\left(1-\frac{16}{m_*}\right)^k-ak\frac{16}{m_*}\left(1-\frac{16}{m_*}\right)^{k-1}\right)\\=&\sum_{k=0}^{(p-1)/2}\frac{\binom{2k}{k}^3}{m_*^k}\left(1-\frac{16}{m_*}\right)^k\left(ak+b-\frac{16}{m_*}ak\frac{m}{m_*}\right)\\=&\sum_{k=0}^{(p-1)/2}\frac{\binom{2k}{k}^3}{m^k}\left(\frac{ak}{16}(m_*-m+32)+b\right)\pmod{p^2}.\end{aligned}$$

On the other hand,

$$\begin{aligned}S=&\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{m_*^k}\sum_{n=k}^{p-1}(an+b)\frac{\binom{2(n-k)}{n-k}^2}{m_*^{n-k}}\\\equiv&\sum_{k=0}^{(p-1)/2}\frac{\binom{2k}{k}^2}{m_*^k}\sum_{j=0}^{(p-1)/2}(a(k+j)+b)\frac{\binom{2j}{j}^2}{m_*^j}\\=&2a\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^2}{m_*^k}\sum_{j=0}^{p-1}\frac{\binom{2j}{j}^2}{m_*^j}+b\left(\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{m_*^k}\right)^2\pmod{p^2}.\end{aligned}$$

Combining the above, we obtain (2.1). □

Lemma 2.2. *Let $p=2n+1$ be an odd prime. Then*

$$P_n(\pm x)\equiv(\pm 1)^n\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{(-16)^k}\left(\frac{x-1}{2}\right)^k \pmod{p^2}. \tag{2.2}$$

Proof. It is well known that $P_n(-x)=(-1)^nP_n(x)$. As noted by van Hammer [vH], for each $k=0,\ldots,n$ we have

$$\begin{aligned}\binom{n}{k}\binom{n+k}{k}=&\binom{n}{k}\binom{-n-1}{k}(-1)^k=\binom{(p-1)/2}{k}\binom{(-p-1)/2}{k}(-1)^k\\ \equiv&\binom{-1/2}{k}^2(-1)^k=\frac{\binom{2k}{k}^2}{(-16)^k} \pmod{p^2}.\end{aligned}$$

So (2.2) holds. □

Lemma 2.3. *Let $d\in\{1,2,3,7\}$ and suppose that p is an odd prime with $(\frac{-d}{p})=-1$. Let $\sqrt{-d}$ be a solution of $z^2+d=0$ in the ring of p-adic integers. Write $p=x^2+dy^2$ with $x,y\in\mathbb{Z}$ such that $\pi=x+y\sqrt{-d}\equiv 0 \pmod p$. Then, for $\bar\pi=x-y\sqrt{-d}$ we have*

$$\bar\pi\equiv 2x-\frac{p}{2x}\equiv-\frac{\sqrt{-d}}{2}\left(4y-\frac{p}{dy}\right) \pmod{p^2}. \tag{2.3}$$

Proof. Note that

$$\bar\pi=2x-(x+y\sqrt{-d})=2x-\frac{p}{x-\sqrt{-d}y}\equiv 2x-\frac{p}{2x} \pmod{p^2}$$

and

$$\bar\pi=-2y\sqrt{-d}+\frac{p}{x-y\sqrt{-d}}\equiv-2y\sqrt{-d}+\frac{p}{-2y\sqrt{-d}}=-\frac{\sqrt{-d}}{2}\left(4y-\frac{p}{dy}\right) \pmod{p^2}.$$

This concludes the proof. □

Lemma 2.4. *Let $p=2n+1$ be an odd prime. Let $d\in\{2,3,7\}$ and suppose that $(\frac{-d}{p})=1$. Write $p=x^2+dy^2$ such that $\pi=x+y\sqrt{-d}\equiv 0 \pmod p$, and that $x\equiv 1 \pmod 4$ if $d=2$, and $x+y\equiv 1 \pmod 4$ if $d\in\{3,7\}$. Set $\bar\pi=x-y\sqrt{-d}$.*

(i) *If $d=2$, then*

$$P_n(\sqrt{2})\equiv i^{-y}\bar\pi \pmod{p^2}.$$

(ii) *If $d=3$, then*

$$P_n(\sqrt{-3})\equiv(-1)^y\bar\pi \pmod{p^2}$$

and

$$P_n\left(\frac{\sqrt{3}}{2}\right)\equiv(-i)^n\bar{\pi}\pmod{p^2}.$$

(iii) *If* $d=7$, *then*

$$P_n(3\sqrt{-7})\equiv(-1)^{n+y}\bar{\pi}\pmod{p^2}$$

and

$$P_n\left(\frac{3\sqrt{7}}{8}\right)\equiv i^n\bar{\pi}\pmod{p^2}.$$

Proof. This follows from [CV, (48)] in the case $m=r=1$ and Table III of [CV, p. 282]. However, we note that certain entries of Table III should be corrected. This results in our addition of $(-1)^y$ in part (ii) and the first congruence in part (iii). (One can check that our correction is necessary via computation.) □

3. Proofs of Theorems 1.1-1.4

Proof of Theorem 1.1. Without loss of generality we assume that $\pi=x+y\sqrt{-2}\equiv 0\pmod p$. (If not, we may use $-y$ instead of y.) It is easy to see that $(-1)^y=(-1)^n$. By Lemmas 2.2 and 2.4,

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{32^k}(1\pm\sqrt{2})^k\equiv(\mp1)^nP_n(\sqrt{2})\equiv(\mp1)^n(-i)^y\bar{\pi}=(\pm1)^ni^y\bar{\pi}\pmod{p^2}. \tag{3.1}$$

Thus

$$\begin{aligned}2\sqrt{2}\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{32^k}P_k=&\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{32^k}((1+\sqrt{2})^k-(1-\sqrt{2})^k)\\ \equiv& i^y\bar{\pi}(1-(-1)^n)\equiv\begin{cases}0\pmod{p^2} & \text{if } p\equiv1\pmod 8,\\ 2i^y\bar{\pi}\pmod{p^2} & \text{if } p\equiv 3\pmod 8.\end{cases}\end{aligned}$$

By Lemma 2.3,

$$\bar{\pi}\equiv-\sqrt{-2}\left(2y-\frac{p}{4y}\right)\pmod{p^2}.$$

So, if $p\equiv3\pmod 8$ then

$$2\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{32^k}P_k\equiv i^y\sqrt{2}\bar{\pi}\equiv(-2i)i^y\left(2y-\frac{p}{4y}\right)=2i^{y-1}\left(2y-\frac{p}{4y}\right)\pmod{p^2}$$

and hence

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{32^k}P_k\equiv(-1)^{(y-1)/2}\left(2y-\frac{p}{4y}\right)\pmod{p^2}$$

as desired. Note that

$$\begin{aligned}\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{32^k}Q_k=&\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{32^k}((1+\sqrt{2})^k+(1-\sqrt{2})^k)\\ \equiv&i^y\bar{\pi}(1+(-1)^n)\equiv\begin{cases}2i^y\bar{\pi}\pmod{p^2}&\text{if }p\equiv1\pmod 8,\\ 0\pmod{p^2}&\text{if }p\equiv3\pmod 8.\end{cases}\end{aligned}$$

If $p\equiv1\pmod 8$, then $2\mid y$ and

$$\frac{y}{2}\equiv\left(\frac{y}{2}\right)^2+\frac{x-1}{4}\cdot\frac{x-1}{2}=\frac{y^2}{4}+\frac{x^2-1}{8}-\frac{x-1}{4}=\frac{p-1}{8}-\frac{x-1}{4}\pmod 2,$$

hence

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{32^k}Q_k\equiv2(-1)^{y/2}\bar{\pi}\equiv(-1)^{(p-1)/8+(x-1)/4}\left(4x-\frac{p}{x}\right)\pmod{p^2}$$

with the help of Lemma 2.3.

Applying Lemma 2.1 with $m=-64$, $m_*=-32\pm32\sqrt{2}$, $a=2$ and $b=2\pm\sqrt{2}$, we obtain that

$$\begin{aligned}&(2\pm\sqrt{2})\sum_{k=0}^{p-1}(4k+1)\frac{\binom{2k}{k}^3}{(-64)^k}\\ \equiv&4\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^3}{32^k}(1\pm\sqrt{2})^k\sum_{j=0}^{p-1}\frac{\binom{2j}{j}^2}{32^j}(1\pm\sqrt{2})^j\\ &+(2\pm\sqrt{2})\left(\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{32^k}(1\pm\sqrt{2})^k\right)^2\pmod{p^2}.\end{aligned}$$

This, together with the van Hammer-Mortenson congruence

$$\sum_{k=0}^{p-1}(4k+1)\frac{\binom{2k}{k}^3}{(-64)^k}\equiv p\left(\frac{-1}{p}\right)\pmod{p^3}$$

(which was conjectured by van Hamme [vH] and proved by E. Mortenson, and further refined by the author [Su6]) and (3.1), yields that

$$\begin{aligned}&\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^2}{32^k}(1\pm\sqrt{2})^k\\ \equiv&\frac{2\pm\sqrt{2}}{4}(\mp1)^ni^y\frac{p(-1)^n-(-1)^y\bar{\pi}^2}{\bar{\pi}}\\ =&\frac{2\pm\sqrt{2}}{4}(\pm1)^ni^y\left(\frac{p}{\bar{\pi}}-\bar{\pi}\right)=\frac{2\pm\sqrt{2}}{2}(\pm1)^ni^yy\sqrt{-2}\pmod{p^2}.\end{aligned}$$

Therefore

$$\begin{aligned}2\sqrt{2}\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^2}{32^k}P_k=&\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^2}{32^k}\left((1+\sqrt{2})^k-(1-\sqrt{2})^k\right)\\ \equiv& i^y y\sqrt{-2}\left(\frac{2+\sqrt{2}}{2}-(-1)^n\frac{2-\sqrt{2}}{2}\right)\pmod{p^2}\end{aligned}$$

and hence

$$\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^2}{32^k}P_k\equiv\begin{cases}i^y y\sqrt{-2}/2\pmod{p^2} & \text{if } p\equiv 1\pmod 8,\\ i^{y+1}y=(-1)^{(y+1)/2}y\pmod{p^2} & \text{if } p\equiv 3\pmod 8.\end{cases}$$

Note that if $p\equiv 1\pmod 8$ then

$$i^y=(-1)^{y/2}=(-1)^{(p-1)/8+(x-1)/4}$$

and

$$y\sqrt{-2}=\frac{p}{x-y\sqrt{-2}}-x\equiv\frac{p}{2x}-x\quad\pmod{p^2}.$$

Observe that

$$\begin{aligned}&\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^2}{32^k}Q_k\\ \equiv& i^y y\sqrt{-2}\left(\frac{2+\sqrt{2}}{2}+(-1)^n\frac{2-\sqrt{2}}{2}\right)\\ \equiv&\begin{cases}i^y y2\sqrt{-2}\equiv 4\sum_{k=0}^{p-1}kP_k\binom{2k}{k}^2/32^k\pmod{p^2} & \text{if } p\equiv 1\pmod 8,\\ i^{y+1}2y=(-1)^{(y+1)/2}2y\pmod{p^2} & \text{if } p\equiv 3\pmod 8.\end{cases}\end{aligned}$$

Combining the above we immediately get all the results stated in Theorem 1.1. □

Proof of Theorem 1.2. Let $\sqrt{-3}$ be a root of $z^2+3=0$ in the ring $\mathbb{Z}_p$ of p-adic integers. Without loss of generality we suppose that $\pi=x+\sqrt{-3}y\equiv 0\pmod p$ and $x+y\equiv 1\pmod 4$. In view of Lemmas 2.2 and 2.4, we have

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{(-16)^k}\left(\frac{-1\pm\sqrt{-3}}{2}\right)^k\equiv(\pm1)^nP_n(\sqrt{-3})\equiv(\pm1)^n(-1)^y\bar{\pi}\quad\pmod{p^2}.$$

By Lemma 2.3,

$$\bar{\pi}\equiv 2x-\frac{p}{2x}\equiv-\frac{\sqrt{-3}}{2}\left(4y-\frac{p}{3y}\right)\quad\pmod{p^2}.$$

Now, applying Lemma 2.1 with $m = 16$, $m_* = 8(1 \pm \sqrt{-3})$, $a = 6$ and $b = 3 \pm \sqrt{-3}$, and noting that

$$\sum_{k=0}^{p-1}(3k+1)\frac{\binom{2k}{k}^3}{16^k} \equiv p \pmod{p^2}$$

by [GZ], we get

$$\begin{aligned}\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^2}{(-16)^k}\left(\frac{-1\pm\sqrt{-3}}{2}\right)^k \equiv& \frac{3\pm\sqrt{-3}}{12}(\pm1)^n(-1)^y\frac{p-\bar{\pi}^2}{\bar{\pi}}\\ =&\frac{3\pm\sqrt{-3}}{6}(\pm1)^n(-1)^y y\sqrt{-3} \pmod{p^2}.\end{aligned}$$

Note that $\omega = (-1+\sqrt{-3})/2$ and $\bar{\omega} = (-1-\sqrt{-3})/2$ are primitive cubic roots of unity. As $\omega + \bar{\omega} = -1$ and $\omega\bar{\omega} = 1$, we see that

$$u_n(-1,1) = \frac{\omega^n - \bar{\omega}^n}{\sqrt{-3}} = \left(\frac{n}{3}\right) \text{ and } v_n(-1,1) = \omega^n + \bar{\omega}^n = 3[3 \mid n] - 1.$$

Combining the above, similar to the proof of Theorem 1.1, we get all the results in Theorem 1.2. □

Proof of Theorem 1.3. Let $\sqrt{-7}$ be a root of $z^2 + 7 = 0$ in the ring $\mathbb{Z}_p$ of p-adic integers. Without loss of generality we suppose that $\pi = x + \sqrt{-7}y \equiv 0 \pmod{p}$ and $x + y \equiv 1 \pmod{4}$. In view of Lemmas 2.2 and 2.4, we have

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}^2}{16^k}\left(\frac{1\pm3\sqrt{-7}}{2}\right)^k \equiv (\mp1)^n P_n(3\sqrt{-7}) \equiv (\pm1)^n(-1)^y\bar{\pi} \pmod{p^2}.$$

By Lemma 2.3,

$$\bar{\pi} \equiv 2x - \frac{p}{2x} \equiv -\frac{\sqrt{-7}}{2}\left(4y - \frac{p}{7y}\right) \pmod{p^2}.$$

Now, applying Lemma 2.1 with $m = 1$, $m_* = (1 \pm 3\sqrt{-7})/2$, $a = 28$ and $b = 21 \pm \sqrt{-7}$, and noting that

$$\sum_{k=0}^{p-1}(21k+8)\binom{2k}{k}^3 \equiv 8p \pmod{p^3}$$

by [Su2], we get

$$\begin{aligned}\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^2}{16^k}\left(\frac{1\pm3\sqrt{-7}}{2}\right)^k \equiv&(\pm1)^n(-1)^y\frac{21\mp\sqrt{-7}}{56}\cdot\frac{p-\bar{\pi}^2}{\bar{\pi}}\\ =&\frac{21\mp\sqrt{-7}}{28}(\pm1)^n(-1)^y y\sqrt{-7} \pmod{p^2}.\end{aligned}$$

Note that

$$y\sqrt{-7}=\frac{p}{x-y\sqrt{-7}}-x\equiv\frac{p}{2x}-x\pmod{p^2}.$$

Also, $(-1)^y=(\frac{2}{p})$ if $p\equiv 1\pmod 4$, and $(-1)^{(y-1)/2}=(\frac{2}{p})$ if $p\equiv 3\pmod 4$. As

$$3\sqrt{-7}u_k(1,16)=\left(\frac{1+3\sqrt{-7}}{2}\right)^k-\left(\frac{1-3\sqrt{-7}}{2}\right)^k$$

and

$$v_k(1,16)=\left(\frac{1+3\sqrt{-7}}{2}\right)^k+\left(\frac{1-3\sqrt{-7}}{2}\right)^k,$$

we immediately obtain the desired results from the above. □

Proof of Theorem 1.4. The proof is similar to the proofs of Theorems 1.1 and 1.2. We apply Lemma 2.1 with $m=256$, $m_*=64/(2\pm\sqrt{3})$, $a=3$ and $b=3\pm 2\sqrt{3}$, and use the congruence

$$\sum_{k=0}^{(p-1)/2}\frac{6k+1}{256^k}\binom{2k}{k}^3\equiv p\left(\frac{-1}{p}\right)\pmod{p^4}$$

conjectured by van Hamme [vH] and confirmed by L. Long [L]. □

3. Proof of Theorem 1.5

Lemma 4.1. *Let $p=2n+1$ be an odd prime. For each $k=0,\ldots,n$ we have*

$$\binom{2n-k}{k}\equiv(-1)^k\binom{2k}{k}(1-p(H_{2k}-H_k))\pmod{p^2}.\qquad(4.1)$$

Proof. Observe that

$$\begin{aligned}(-1)^k\binom{2n-k}{k}&=(-1)^k\prod_{0<j\leqslant k}\frac{p-k-j}{j}=\prod_{0<j\leqslant k}\frac{j+k-p}{j}\\&=\prod_{0<j\leqslant k}\frac{j+k}{j}\times\prod_{0<j\leqslant k}\left(1-\frac{p}{j+k}\right)\\&\equiv\binom{2k}{k}\left(1-\sum_{0<j\leqslant k}\frac{p}{j+k}\right)=\binom{2k}{k}(1-p(H_{2k}-H_k))\pmod{p^2}.\end{aligned}$$

This concludes the proof. □

Theorem 4.1. *Let $p = 2n + 1$ be an odd prime and let m be an integer not divisible by p. Let $h \in \mathbb{Z}^+$ and $P(x) \in \mathbb{Z}[x]$. Then*

$$\begin{aligned}&\left(\frac{(-1)^h m}{p}\right)\sum_{k=0}^{n} P(k)\frac{\binom{2k}{k}^h}{m^k}\\ &\equiv \sum_{k=0}^{n}\frac{\binom{2k}{k}^h}{\bar{m}^k}\left(\left(1+\frac{p}{2}q_p(\bar{m})\right)P\left(-k-\frac{1}{2}\right)+\frac{p}{2}P'\left(-k-\frac{1}{2}\right)\right)\\ &\quad - ph\sum_{k=0}^{n}\frac{\binom{2k}{k}^h}{\bar{m}^k}\left(P\left(-k-\frac{1}{2}\right)(H_{2k}-H_k)\right) \pmod{p^2},\end{aligned}$$

where $\bar{m} = 16^h/m$ and $q_p(\bar{m}) = (\bar{m}^{p-1}-1)/p$.

Proof. Since

$$\binom{n+k}{n-k} = \binom{n+k}{2k} \equiv \frac{\binom{2k}{k}}{(-16)^k} \pmod{p^2} \quad \text{for all } k = 0, \dots, n,$$

we have

$$\begin{aligned}\sum_{k=0}^{n} P(k)\frac{\binom{2k}{k}^h}{m^k} &\equiv \sum_{k=0}^{n} P(k)\binom{n+k}{n-k}^h\frac{(-16)^{hk}}{m^k}\\ &= \sum_{k=0}^{n} P(n-k)\binom{2n-k}{k}^h((-1)^h\bar{m})^{n-k} \pmod{p^2}.\end{aligned}$$

So, with the help of Lemma 4.1, we get

$$\sum_{k=0}^{n} P(k)\frac{\binom{2k}{k}^h}{m^k} \equiv \sum_{k=0}^{n} P(n-k)\frac{\binom{2k}{k}^h}{\bar{m}^k}(1-p(H_{2k}-H_k))^h((-1)^h\bar{m})^n \pmod{p^2}. \tag{4.2}$$

Since

$$\left(\frac{p}{2}+x\right)^j - x^j \equiv j\frac{p}{2}x^{j-1} \equiv \frac{p}{2}\frac{d}{dx}x^j \pmod{p^2}$$

for all $j = 0, 1, 2, \dots$, we see that

$$P\left(\frac{p}{2}+x\right) - P(x) \equiv \frac{p}{2}P'(x) \pmod{p^2}.$$

Thus

$$P(n-k) = P\left(\frac{p}{2}-k-\frac{1}{2}\right) \equiv P\left(-k-\frac{1}{2}\right) + \frac{p}{2}P'\left(-k-\frac{1}{2}\right) \pmod{p^2}$$

for any $k = 0, 1, \ldots, n$. So, from (4.2) we obtain

$$\begin{aligned}&\left(\frac{(-1)^h}{p}\right)\sum_{k=0}^{n} P(k)\frac{\binom{2k}{k}^h}{m^k}\\ \equiv&\sum_{k=0}^{n}\left(P\left(-k-\frac{1}{2}\right)+\frac{p}{2}P'\left(-k-\frac{1}{2}\right)\right)\frac{\binom{2k}{k}^h}{\bar{m}^k}(1-ph(H_{2k}-H_k))\bar{m}^n \pmod{p^2}.\end{aligned} \tag{4.3}$$

Since

$$\bar{m}^n = \frac{4^{(p-1)h}}{m^n} \equiv \left(\frac{m}{p}\right) \pmod{p},$$

we have

$$\begin{aligned}\bar{m}^{p-1}-1=&\left(\bar{m}^n+\left(\frac{m}{p}\right)\right)\left(\bar{m}^n-\left(\frac{m}{p}\right)\right)\\ \equiv&2\left(\frac{m}{p}\right)\left(\bar{m}^n-\left(\frac{m}{p}\right)\right) \pmod{p^2}\end{aligned}$$

and hence

$$1+\frac{p}{2}q_p(\bar{m}) \equiv \left(\frac{m}{p}\right)\bar{m}^n \pmod{p^2}.$$

Combining this with (4.3) we immediately obtain the desired result. □

Proof of Theorem 1.5. Let $n = (p-1)/2$. Note that $p \mid \binom{2k}{k}$ for all $k = n+1, \ldots, p-1$. Applying Theorem 4.1 with $h = 3$ and $P(x) = ax + b$ we then get the desired result.

Below we give some corollaries of Theorem 1.5.

Corollary 4.1. *Let $p > 3$ be a prime. Then*

$$\begin{aligned}&\left(\frac{-1}{p}\right)\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^3}{(-64)^k}(H_{2k}-H_k)-\frac{1}{6}\\ \equiv&\begin{cases}-(3q_p(2)+2)x^2/3 \pmod{p} & \text{if } p=x^2+2y^2,\\ 0 \pmod{p} & \text{if } p\equiv 5,7 \pmod{8}.\end{cases}\end{aligned} \tag{4.4}$$

When $p \equiv 1 \pmod{4}$ and $p = x^2 + y^2$ with x odd and y even, we have

$$\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^3}{64^k}(H_{2k}-H_k) \equiv -\frac{x^2}{3}(3q_p(2)+2) \pmod{p}. \tag{4.5}$$

Proof. Recall the known congruence

$$\sum_{k=0}^{p-1}\frac{4k+1}{(-64)^k}\binom{2k}{k}^3 \equiv p\left(\frac{-1}{p}\right) \pmod{p^3}$$

due van Hamme [vH] and Mortenson. In view of this and (1.3) and (1.9), Theorem 1.1 with $m=-64$, $a=4$ and $b=1$ yields (4.4).

Let $p\equiv 1\pmod 4$ and write $p=x^2+y^2$ with x odd and y even. The author [Su2, Conj. 5.9] conjectured that

$$\sum_{k=0}^{p-1}\frac{4k+1}{64^k}\binom{2k}{k}^3\equiv 0\pmod{p^2},$$

and this was confirmed by Z.-H. Sun [S2]. In view of this and (1.0) and (1.9), we obtain (4.5) from Theorem 1.1 with $m=64$, $a=4$ and $b=1$.

Corollary 4.2. *Let $p>3$ be a prime. Then*

$$\begin{aligned}&\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^3}{16^k}(H_{2k}-H_k)-\frac16\\ \equiv&\left(\frac{-1}{p}\right)\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^3}{256^k}(H_{2k}-H_k)-\frac16\\ \equiv&\begin{cases}-2x^2(4q_p(2)+3)/9\pmod p & \text{if } p=x^2+3y^2,\\ 0\pmod p & \text{if } p\equiv 2\pmod 3.\end{cases}\end{aligned}\tag{4.6}$$

Proof. van Hamme [vH] conjectured the congruence

$$\sum_{k=0}^{(p-1)/2}\frac{6k+1}{256^k}\binom{2k}{k}^3\equiv p\left(\frac{-1}{p}\right)\pmod{p^4}$$

and this was confirmed by L. Long [L]. The author [Su2] conjectured

$$\sum_{k=0}^{(p-1)/2}\frac{3k+1}{16^k}\binom{2k}{k}^3\equiv p+2\left(\frac{-1}{p}\right)p^3E_{p-3}\pmod{p^4}$$

(where $E_0,E_1,\ldots$ are Euler numbers), and its mod p^3 was proved in [GZ]. In view of this and (1.4)-(1.5) and (1.9), we deduce (4.6) by applying Theorem 1.1 with $(a,b,m)=(6,1,256),(3,1,16)$. □

van Hammer [vH] conjectured that for any prime $p>3$ we have

$$\sum_{k=0}^{(p-1)/2}\frac{42k+5}{4096^k}\binom{2k}{k}^3\equiv 5p\left(\frac{-1}{p}\right)\pmod{p^4}.$$

This is even open modulo p^2.

Corollary 4.3. *Let $p>3$ be a prime. Then*

$$\begin{aligned}&\sum_{k=0}^{p-1}\frac{42k+5}{4096^k}\binom{2k}{k}^3\equiv 5p\left(\frac{-1}{p}\right)\pmod{p^2}\\ \iff&\sum_{k=0}^{p-1}k\binom{2k}{k}^3(H_{2k}-H_k)-\frac{1}{6}\\ &\equiv\begin{cases}-2x^2/3\pmod p & \text{if } (\frac{p}{7})=1\ \&\ p=x^2+7y^2,\\ 0\pmod p & \text{if } (\frac{p}{7})=-1.\end{cases}\\ \iff&\left(\frac{-1}{p}\right)\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^3}{4096^k}(H_{2k}-H_k)-\frac{1}{6}\\ &\equiv\begin{cases}-2(10q_p(2)+7)x^2/21\pmod p & \text{if } (\frac{p}{7})=1\ \&\ p=x^2+7y^2,\\ 0\pmod p & \text{if } (\frac{p}{7})=-1.\end{cases}\end{aligned}$$

Proof. By [Su2, Theorem 3] we have

$$\sum_{k=0}^{p-1}(21k+8)\binom{2k}{k}^3\equiv 8p\pmod{p^4}.$$

Applying Theorem 1.1 with $(a,b,m)=(21,8,1),(42,5,4096)$ and noting (1.1),(1.6) and (1.9) we obtain the desired result. □

van Hammer [vH] conjectured that for any prime $p>3$ we have

$$\sum_{k=0}^{(p-1)/2}\frac{6k+1}{(-512)^k}\binom{2k}{k}^3\equiv p\left(\frac{-2}{p}\right)\pmod{p^3}.$$

This is even open modulo p^2.

Corollary 4.4. *Let $p>3$ be a prime. Then*

$$\begin{aligned}&\sum_{k=0}^{p-1}\frac{6k+1}{(-512)^k}\binom{2k}{k}^3\equiv p\left(\frac{-2}{p}\right)\pmod{p^2}\\ \iff&\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^3}{(-8)^k}(H_{2k}-H_k)-\frac{(-1)^{(p-1)/2}}{6}\\ &\equiv\begin{cases}-2(q_p(2)+1)x^2/3\pmod p & \text{if } p=x^2+y^2\ (2\nmid x),\\ 0\pmod p & \text{if } p\equiv 3\pmod 4.\end{cases}\\ \iff&\left(\frac{-2}{p}\right)\sum_{k=0}^{p-1}\frac{k\binom{2k}{k}^3}{(-512)^k}(H_{2k}-H_k)-\frac{1}{6}\\ &\equiv\begin{cases}-(3q_p(2)+2)x^2/3\pmod p & \text{if } p=x^2+y^2\ (2\nmid x),\\ 0\pmod p & \text{if } p\equiv 3\pmod 4.\end{cases}\end{aligned}$$

Proof. The author [Su2] conjectured that

$$\sum_{k=0}^{p-1} \frac{3k+1}{(-8)^k} \equiv p\left(\frac{-1}{p}\right) + p^3 E_{p-3} \pmod{p^4}.$$

and this was proved in [GZ] mod p^3. Applying Theorem 1.4 with $(a,b,m) = (6,1,-512),(3,1,-8)$ and noting (1.2) and (1.9) we obtain the desired result. □

Our following conjecture seems difficult.

Conjecture 4.1. *Let p be an odd prime.*

(i) *If $p \equiv 1 \pmod 4$ then*

$$\begin{aligned}\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^3}{(-8)^k} \sum_{k<j\leqslant 2k} \frac{1}{j} \equiv& \frac{1}{2}\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^3}{64^k} \sum_{k<j\leqslant 2k} \frac{1}{j}\\ \equiv& \frac{1}{3}\left(\frac{2}{p}\right)\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^3}{(-512)^k} \sum_{k<j\leqslant 2k} \frac{1}{j} \pmod{p^2};\end{aligned}$$

when $p \equiv 3 \pmod 4$ we have

$$\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^3}{(-8)^k} \sum_{k<j\leqslant 2k} \frac{1}{j} \equiv -\frac{7}{2}\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^3}{64^k} \sum_{k<j\leqslant 2k} \frac{1}{j} \pmod{p^2},$$

$$\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^3}{64^k} \sum_{k<j\leqslant 2k} \frac{1}{j} \equiv -\left(\frac{2}{p}\right)\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^3}{(-512)^k} \sum_{k<j\leqslant 2k} \frac{1}{j} \pmod{p^2},$$

(ii) *If $p \equiv 1 \pmod 3$ then*

$$\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^3}{16^k} \sum_{k<j\leqslant 2k} \frac{1}{j} \equiv \frac{1}{2}\left(\frac{-1}{p}\right)\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^3}{256^k} \sum_{k<j\leqslant 2k} \frac{1}{j} \pmod{p^2}.$$

If $p \equiv 2 \pmod 3$ then

$$\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^3}{256^k} \sum_{k<j\leqslant 2k} \frac{1}{j} \equiv 0 \pmod{p^2}.$$

(iii) *If $p > 3$ and $p \equiv 3,5,6 \pmod 7$, then*

$$\sum_{k=0}^{(p-1)/2} \binom{2k}{k}^3 \sum_{k<j\leqslant 2k} \frac{1}{j} \equiv 0 \pmod{p^2}.$$

References

[A] S. Ahlgren, *Gaussian hypergeometric series and combinatorial congruences*, in: Symbolic Computation, Number Theory, Special Functions, Physics and Combinatorics (Gainesville, FI, 1999), pp. 1-12, Dev. Math., Vol. 4, Kluwer, Dordrecht, 2001.

[BEW] B. C. Berndt, R. J. Evans and K. S. Williams, *Gauss and Jacobi Sums*, John Wiley & Sons, 1998.

[CDE] S. Chowla, B. Dwork and R. J. Evans, *On the mod* p^2 *determination of* $\binom{(p-1)/2}{(p-1)/4}$, J. Number Theory **24** (1986), 188–196.

[CV] M. J. Coster and L. van Hamme, *Supercongruences of Atkin and Swinnerton-Dyer type for Legendre polynomials*, J. Number Theory **38** (1991), 265–286.

[C] D. A. Cox, *Primes of the Form* $x^2 + ny^2$, John Wiley & Sons, 1989.

[GZ] J. Guillera and W. Zudilin, *"Divergent" Ramanujan-type supercongruences*, Proc. Amer. Math. Soc. **140** (2012), 765–777.

[HW84] R. H. Hudson and K. S. Williams, *Binomial coefficients and Jacobi sums*, Trans. Amer. Math. Soc. **281** (1984), 431–505.

[JO] M. Jameson and K. Ono, *Personal communication via e-mail*, March 2010.

[KLMSY] J. Kibelbek, L. Long, K. Moss, B. Sheller and H. Yuan, *Supercongruences and complex multiplication*, preprint, arXiv:1210.4489v1 (Oct. 16, 2012).

[L] L. Long, *Hypergeometric evaluation identities and supercongruences*, Pacific J. Math. **249** (2011), 405–418.

[S1] Z. H. Sun, *Congruences concerning Legendre polynomials*, Proc. Amer. Math. Soc. **139** (2011), 1915–1929.

[S2] Z. H. Sun, *Congruences concerning Legendre polynomials II*, preprint, arXiv:1012.3898v2 (August 3, 2012).

[Su] Z. W. Sun, *Open conjectures on congruences*, preprint, arXiv:0911.5665.

[Su1] Z. W. Sun, *On congruences related to central binomial coefficients*, J. Number Theory **131** (2011), 2219–2238.

[Su2] Z. W. Sun, *Super congruences and Euler numbers*, Sci. China Math. **54** (2011), 2509-2535.

[Su3] Z. W. Sun, *On sums of Apéry polynomials and related congruences*, J. Number Theory **132** (2012), 2673–2699.

[Su4] Z. W. Sun, *On sums of binomial coefficients modulo* p^2, Colloq. Math. **127** (2012), 39–54.

[Su5] Z. W. Sun, *On sums involving products of three binomial coefficients*, Acta Arith. **in press.** `arXiv:1012.3141`.

[Su6] Z. W. Sun, *A refinement of a congruence result by van Hamme and Mortenson*, Illinois J. Math., in press. arXiv:1011.1902.

[Su7] Z. W. Sun, *On sums of central binomial and trinomial coefficients*, in: Number Theory–New York Seminar (ed., M. B. Nathanson), Springer, in press.

[Su8] Z. W. Sun, *Conjectures and results on* x^2 *mod* p^2 *with* $4p = x^2 + dy^2$, in: Number Theory and the Related Topics (eds., Y. Ouyang, C. Xing, F. Xu and P. Zhang), Higher Education Press & International Press, Beijing and Boston, in press.

[vH] L. van Hamme, *Some conjectures concerning partial sums of generalized hypergeometric series*, in: p-adic Functional Analysis (Nijmegen, 1996), pp. 223–236, Lecture Notes in Pure and Appl. Math., Vol. 192, Dekker, 1997.